

NGC 7582: The Prototype Narrow-Line X-ray Galaxy

Jonathan F. Schachter, Fabrizio Fiore¹, Martin Elvis, Smita Mathur

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

Andrew S. Wilson

Astronomy Department, University of Maryland, College Park MD 21218

Jon A. Morse

Center for Astrophysics and Space Astronomy, University of Colorado, Campus Box 389, Boulder, CO 80309

Hisamitsu Awaki

Department of Physics, Faculty of Science, Kyoto University, Kitashirakawa, Sakyo, Kyoto, 606-01, Japan

Kazushi Iwasawa

Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, United Kingdom

Accepted for publication in The Astrophysical Journal (Letters)

¹present address: Osservatorio Astronomico di Roma, Via dell' Osservatorio, I-00140 Monteporzio, Italy

ABSTRACT

NGC 7582 is a candidate prototype of the Narrow Line X-ray Galaxies (NLXGs) found in deep X-ray surveys. An ASCA observation shows the hard (>3 keV) X-ray continuum of NGC 7582 drops 40% in ~ 6 ks, implying an AGN, while the soft band (< 3 keV) does not drop in concert with the hard continuum, requiring a separate component. The X-ray spectrum of NGC 7582 also shows a clear 0.5–2 keV soft ($kT = 0.8^{+0.9}_{-0.3}$ keV or $\Gamma = 2.4 \pm 0.6$; $L_X = 6 \times 10^{40}$ ergs s^{-1}) low-energy component, in addition to a heavily absorbed [$N_H = (6 \pm 2) \times 10^{22}$ cm^{-2}] and variable 2–10 keV power law [$\Gamma = 0.7^{+0.3}_{-0.4}$; $L_X = (1.7 - 2.3) \times 10^{42}$ ergs s^{-1}]. This is one of the flattest 2–10 keV slopes in any AGN observed with ASCA. (The ROSAT HRI image of NGC 7582 further suggests extent to the SE.)

These observations make it clear that the hard X-ray emission of NGC 7582, the most ‘narrow-line’ of the NLXGs, is associated with an AGN. The strong suggestion is that *all* NLXGs are obscured AGNs, as hypothesized to explain the X-ray background spectral paradox. The separate soft X-ray component makes NGC 7582 (and by extension other NLXGs) detectable as a ROSAT source.

Subject headings: galaxies: active, galaxies: Seyfert, X-rays: galaxies, BL Lacertae objects: general

1. Introduction

Deep X-ray surveys with ROSAT have revealed a rapidly increasing number of ‘Narrow-Line X-ray Galaxies’ (NLXGs) at moderate redshifts ($z \sim 0.1 - 0.5$) and luminosities ($L_X \sim 10^{42} - 10^{43}$ ergs s $^{-1}$, Boyle et al. 1995, Georgantopoulos et al. 1995). NLXGs were first discovered in early 2–10 keV sky surveys (e.g., *Ariel V*, Ward et al. 1978; HEAO-1, Piccinotti et al. 1982). Since these NLXGs are highly absorbed ($N_H \sim 10^{22-23.5}$ cm $^{-2}$; Turner & Pounds 1989) at low X-ray energies, their ROSAT counterparts may be the absorbed AGN population posited (Comastri et al. 1995) to solve both the hard X-ray background ‘spectral paradox’ (Boldt 1987) and the hard X-ray/soft X-ray source counts discrepancy (Gendreau et al. 1995b, Georgantopoulos, et al., 1997). This picture is not without problems however: it predicts extremely hard ROSAT spectra, and ASCA fluxes many times brighter than ROSAT. Yet neither is found (Georgantopoulos, et al., 1997, Romero-Colmenero et al., 1996).

The NLXGs may not all be AGN; in high quality optical spectra half the NLXGs have no broad H α (Boyle et al. 1995) and so may be starbursts, rather than AGN. The brightest NLXG is NGC 7582, which has linewidths < 200 km s $^{-1}$ (Morris et al. 1985; Wilson & Nath 1990) and no broad components out to Brackett- γ (Moorwood & Oliva 1990). As such, NGC 7582 is the best and the brightest NLXG candidate for being starburst dominated, and may serve as a prototype of the new faint NLXG population. X-ray observations can be critical to determining the presence of an AGN.

Previous X-ray spectra (Mushotzky 1982, Warwick et al. 1993, Turner & Pounds 1989) and variability measurements (Mushotzky 1982) of NGC 7582 however, have been hampered by bright confusing sources: the background cluster Abell S1101 is 53’ away (Charles & Phillips 1982), while the BL Lac object PKS 2316–423

is only 17’ away (Crawford & Fabian 1994; Stocke et al. 1991).

In this *Letter* we describe ASCA SIS and ROSAT HRI observations of NGC 7582, separating the contributions of the AGN and the BL Lac. These demonstrate that even this purely narrow-line object clearly harbors an AGN; simultaneously these data suggest an answer to the problems of the soft X-ray flux from NLXGs.

2. Observations

2.1. ASCA SIS Data

An ASCA observation in SIS 2 CCD mode (along the diagonal) and carefully oriented to place both NGC 7582 and PKS 2316–423 near the axis, was performed on 1994 November 11, acquiring 17.0 ks in SIS0 and 18.2 ks in SIS1 on NGC 7582. Both sources were clearly detected. At 0.1 cts s $^{-1}$, NGC 7582 dominates the hard (2–10 keV) band by a factor of ~ 2.3 . A total of 4067 net SIS counts were detected from NGC 7582, allowing complex spectral fits. We used the `ascascreen` default criteria ² for data screening, extracted the SIS spectra from the recommended 4’ radius region (ABC Guide) and binned to give >10 counts per bin. There were no high background intervals in the SIS full-field light curves.

Soft (0.5–2 keV) and hard (3–8 keV) SIS X-ray light curves (Figure 1) in 8 bins, one per orbit, spanning a 40 ks time base, for the combined SIS0 and SIS1 data, detect NGC 7582 at $> 4\sigma$ in each orbit in both bands. The hard count rate is significantly variable ($\chi^2=21.9$), showing a clear (9.2σ) 40% decrease in ~ 6 ks between bins 5 and 6, and remaining low for the rest of the observa-

²See definitions in Table 5.5.1 of ABC Guide to ASCA Analysis; hereafter, ABC Guide. Maximum angular deviation = 0.01; ELV >10 ; BR_EARTH >20 COR_MIN = 6; PIXL rejection threshold = 75 (standard for 2-CCD mode); remove hot and flickering pixels, event grades 0, 2, 3, and 4. Data from bright and faint (“bright2”) mode, and high- and medium-bit rate data, were combined.

tion³ The soft count rate shows no such drop and is consistent with a constant value ($\chi^2 = 1.3$), excluding the fifth bin, which lies $\sim 5\sigma$ from the mean and may be due to bad background subtraction. Including the 5th bin gives $\chi^2 = 4.28$. The hardness ratio, which is essentially constant through bin 5, is significantly lower in bins 6-8. We refer to the first five bins as the *high state*, and to the remainder as the *low state*.




Fig. 1.— Background-subtracted hard (3-8 keV, top), soft (0.5-2 keV, middle) and hardness ratio (bottom) light curves for NGC 7582 (SIS0 and SIS1 data combined).

We fit the high and low state spectra separately using XSPEC 10.00 (Arnaud 1996), using the 1994 November 9 calibrations. The Galactic line-of-sight N_H is low ($1.48 \times 10^{20} \text{ cm}^{-2}$; Elvis, Wilkes, & Lockman 1989), due to the proximity to the South Galactic pole. Both SIS detectors gave consistent results, and so we report only simultaneous fits (Table 1). A simple power-law fit is unacceptable ($\chi^2_\nu > 1.6$; Model A in Table 1). See Figure 2. (A fit to just the 2-10 keV

³In the PKS 2316-423 SIS data, we find no analogous abrupt change in flux; qualitatively, the BL Lac spectrum is more short-time variable than that of NGC 7582.

range does give an acceptable power-law fit of $\Gamma = 1.4 \pm 0.3$.)




Fig. 2.— Best-fit model, data and χ^2 for the simple-power law fit (Model A) for simultaneous SIS0 (squares) and SIS1 (triangles) high-state observations of NGC 7582.

Addition of a low-energy unabsorbed continuum component, either thermal bremsstrahlung ($kT \sim 0.8 \text{ keV}$) or a second, steep power-law ($\Gamma \sim 2.5$) produces a good fit (Models B and C).

Although a partial covering model (Model D) does significantly better than a single power law (and the underlying power law photon index is more typical of canonical AGN values), adding a soft, unabsorbed component is the preferred model, since an F -test gives a $>99\%$ improvement in the fit going from Model D to Model C. Interestingly, the partial covering model yields emission-line-like residuals near 1 keV, as have been reported in NGC 4151 (Weaver et al. 1994). No changes in the parameters, except normalization, are seen (at 90% confidence) between the two states. The steep power-law needed at low energies is inconsistent with the higher energy power-law at high confidence.

Using the power-law plus bremsstrahlung fit

(Model B), the high (low)-state luminosity, L_X , of NGC 7582 in units of $10^{41} \text{ ergs s}^{-1}$ is 23_{-4}^{+6} (17_{-3}^{+7}) in the 2–10 keV band, and $0.62_{-0.23}^{+0.14}$ ($0.59_{-0.05}^{+0.02}$) in the 0.5–2.0 keV band. So the low energy, ROSAT band, luminosity is 3% – 4% of the 2–10 keV luminosity. (Luminosities are as observed, i.e. uncorrected for absorption. We adopt a distance of 27 Mpc [de Vaucouleurs et al. 1991].)

Absorbing material produces fluorescence lines, notably Fe-K α A narrow ($\sigma=100$ eV) Gaussian line at the cold Fe-K fluorescence line (6.4 keV), plus a power-law in the energy range 3.5–8 keV, gives 90% confidence upper limits on the line equivalent width, <230 eV (SIS1, low state), <310 eV (SIS0, high state). The absorbing N_H would produce a neutral Fe-K edge (at 7.1 keV) with $\tau_{FeK} = 0.2$. Our 90% confidence limit is $\tau_{FeK} \leq 0.2$.

2.2. ROSAT HRI Data

The ROSAT HRI observed NGC 7582 for 17.2 ksec on 1995 May 23, some 6 months after the ASCA observation. The image was *OBI-shifted* to reduce aspect problems (Morse 1994) and smoothed with a $5''$ Gaussian.

A source is found positionally coincident with NGC 7582 (using IRAF/PROS tasks *xexamine*, *imcnts* and a radius= $8''$ circle), containing 225 ± 32 cts.

The source confusion that has dogged NGC 7582 continues with the discovery of a new bright HRI source 2' to the NE (J2000.0: $23^h 18^m 29.9^s$, $-42^\circ 20' 41'' 4$), which has 90 ± 26 counts, and so 40% of the flux of NGC 7582 in the 0.5–2.0 keV band⁴. Assuming a $kT = 1$ keV bremsstrahlung with Galactic absorption, this implies a total luminosity (NGC 7582 + NE source) of $L_X =$

$3.9(\pm 0.5) \times 10^{40} \text{ ergs s}^{-1}$. For the ASCA SIS0 best-fit bremsstrahlung model, the corresponding result is $L_X = 4.9 \times 10^{40} \text{ ergs s}^{-1}$.

The observed full-width half maximum (FWHM) of NGC 7582 (from the IRAF task *imexamine*) is $4.7''$, and of the NE source $5.0''$. Since the expected FWHM for a point source is $6''$ (David et al. 1996), both NGC 7582 and the NE source are unresolved by this measure. However, the HRI image of NGC 7582 appears more asymmetric than that of the NE source. To investigate, we divided the sources into 4 quadrants (Elvis et al., 1983) of radius $20''$, with the dividing lines (1) parallel to the N-S and E-W axes; and (2) rotated by 45° . In both cases, the image of the NE source was found to be consistent with a symmetric profile, but the image of NGC 7582 is asymmetric – brighter on the E side in Case 2 at $> 3 \sigma$, and $> 2.5 \sigma$ fainter on the S side; fainter on the SW side at $> 2.5 \sigma$, in Case 1. Subtracting a scaled and shifted image of the NE source from NGC 7582 gives 2–3 σ residuals to the SE, confirming the asymmetry. However, given the poor S/N in the NE source we cannot rule out the asymmetry in NGC 7582 being an artifact of residual ROSAT aspect solution errors.

3. Discussion

An AGN dominates the hard X-ray emission from NGC 7582. The ASCA hard X-ray flux drops by 40% in ~ 6 ks, implying a source size of $\sim 2 \times 10^{14}$ cm, for which an AGN is the only known possibility. This is 30 times more variable than the X-ray flare previously reported by Mushotzky 1982.

The low energy component in the NGC 7582 X-ray spectrum solves two problems with the Comastri et al. 1995 ‘obscured AGN’ model for the X-ray background: (1) the predicted ROSAT band spectra of NLXGs should be extremely hard, since the shape will be dominated by the strongly energy dependent ($E^{-8/3}$) photoelectric absorption cross-sections (Tucker 1975), yet

⁴For the NE source, the magnitudes of the two stellar objects on the digitized sky survey inside the $4.2''$ radius 2σ HRI error circle, B=20.2 and 21.1 (Anglo-Australian Observatory 1997), allow an AGN or BL Lac identification or, for the brighter optical object, a star of type M0V or later (Schachter et al. 1996).

Romero-Colmenero et al., 1996 find quite normal $\Gamma=1.5$ slopes and Ciliegi, et al., 1997 find an even steeper $\Gamma \sim 2.4$; (2) with the predicted N_H distribution, most (70% – 90%) NLXGs should be *at least* 10 times brighter in ASCA, yet the four ROSAT NLXGs studied by Georgantopoulos, et al., 1997, are just 4–12 times brighter in ASCA, implying a maximum column density of $2.5 \times 10^{22} \text{ cm}^{-2}$ (for $\Gamma=0.8$).

Both of these difficulties are removed by an extra, soft, X-ray spectral feature in NGC 7582: (1) The low energy component has a power-law slope comparable to that of bright ROSAT NLXGs (Ciliegi, et al., 1997); (2) the low energy component prevents the ASCA to ROSAT flux ratio from exceeding 20, more in line with the ratios implied in Georgantopoulos, et al., 1997. Even so a bright source with this spectrum at $1 \times 10^{-13} \text{ ergs cm}^{-2} \text{ s}^{-1}$ in the 2-10 keV band, would be one of the faintest ROSAT sources in the 0.5-2.0 keV band, at $3 \times 10^{-15} \text{ ergs cm}^{-2} \text{ s}^{-1}$. Without the extra soft component, a simple absorbed power-law spectrum would have a flux of $2 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1}$ in ROSAT, and so would be undetectable. The Georgantopoulos, et al., 1997 ASCA/ROSAT ratios for NLXGs indicate somewhat larger uncovered fractions, $\sim 15\%$.

What is this soft X-ray component in the ASCA spectrum? The constancy of the soft X-rays rules out a simple uncovered fraction of the central continuum, and instead (with the possible HRI asymmetries) suggests an extended origin. (Although the hint of soft-band variability should be borne in mind.) Certainly, kiloparsec scale extended soft X-ray emission is known in other AGN (Elvis et al. 1990, Wilson et al. 1992, Morse et al. 1995), where it may be of nuclear or starburst origin (Wilson et al. 1996). Hot gas is the likely dominant emitter in either case. In NGC 7582, emission line ratios (Ward et al. 1980; Baldwin, Phillips, & Terlevich 1981) and widths (Whittle 1992) allow either a type2 Seyfert or an H II region dominated object. Wilson 1988 concludes that the extended ‘diffuse’ radio emis-

sion, the $4''$ radius ($1'' = 130 \text{ pc}$ at 27 Mpc) H α disk (Morris et al. 1985), and the IRAS far-infrared emission in NGC 7582 are all powered by stellar processes, not by an AGN. A starburst/AGN composite seems to be required (Morris et al. 1985; Wilson 1988), especially given the hard X-ray variability.

The X-ray luminosity of the soft component in NGC 7582 is modest, 10–100 times smaller than that of the brighter, reliably identified (ha98) NLXGs from the “CRSS” (Ciliegi, et al., 1997, Romero-Colmenero et al., 1996) and comparable with starburst galaxies (Fabbiano 1989). In order to explain NLXG luminosities up to $10^{43} \text{ erg s}^{-1}$, and to maintain relatively constant soft/hard band X-ray luminosities, it is more plausible that the soft component is a direct result of nuclear activity, rather than indirect via a starburst (Fabian et al. 1998). This favors Compton-scattered nuclear emission, or jet/outflow-related emission over a starburst. Compton reflection could allow some variability if the ‘mirror’ is close to the nuclear source.

The column density within NGC 7582 decreased by a factor ~ 6 in the 6 years prior to the ASCA observations. The best non-imaging X-ray spectrum of NGC 7582 was obtained with Ginga in 1988 October; Warwick et al. 1993 found $N_H \approx 50 \times 10^{22} \text{ cm}^{-2}$, keeping Γ fixed at 1.7. We find, in the ASCA energy range and assuming the same fixed Γ , $N_H = (7.1^{+0.8}_{-0.4}) \times 10^{22} \text{ cm}^{-2}$.⁵ A column density at the Ginga level in NLXGs would render the central AGN component fainter by a factor 10^6 . This is one of the larger variations in column density known in AGN, similar in both timescale and amplitude to those in NGC 4151 (Yaqoob et al., 1993). Changes on this timescale can be created by motions of molecular clouds in a starburst model, but the amplitude may be hard to reproduce.

⁵Neither Abell S1101, which was allowed for in the Warwick et al., analysis, nor PKS 2316–423, which was not, has such a heavily absorbed spectrum (Schachter et al. 1998)

4. Conclusions

NGC 7582 is the brightest example of the class of NLXGs found in the deepest ROSAT observations (Boyle et al. 1995), with no broad ($> 1000 \text{ km s}^{-1}$) components on their optical emission lines, and may serve as the prototype of the class. Our analysis of ASCA and ROSAT observations of NGC 7582 has shown:

1. A rapid ($\sim 6 \text{ ksec}$) 40% decrease in the ASCA SIS hard ($> 3 \text{ keV}$) X-ray flux, requiring a compact, AGN source, with an observed (absorbed) 2–10 keV luminosity of $\sim 2 \times 10^{42} \text{ ergs s}^{-1}$.
2. An additional, separate, emission component is present in the ASCA SIS soft band (0.5–2 keV, $L_X \sim 6 \times 10^{40} \text{ ergs s}^{-1}$) with a steep ($\Gamma=2.4$) unabsorbed spectrum.
3. A heavily obscured ($N_H \approx 6 \times 10^{22} \text{ cm}^{-2}$) nuclear source, that has likely become 6 times less obscured over the last 6 years. This may cause difficulty explaining the absorbers as being associated with nuclear starbursts, as proposed by Fabian et al. 1998.

We also note that the source confusion issues which have dogged NGC 7582 now seem to be resolved:

4. NGC 7582 is the dominant source (by a factor of 2–3) of 2–10 keV emission in the field containing itself and the BL Lac object PKS 2316–423.
5. In the soft X-ray band NGC 7582 dominates the newly discovered source 2 'NE by a factor 2.

These observations make it clear that the hard X-ray emission of NGC 7582, the most 'narrow-line' of the NLXGs, is associated with an AGN. The strong suggestion is that *all* NLXG are obscured AGNs, as hypothesized to explain the

hard X-ray background spectral paradox; but that it is a separate soft X-ray component that makes NLXGs detectable as ROSAT sources, and provides their AGN-like X-ray slope. Quite likely this soft component is extended, and thermal in nature.

We thank the anonymous referee for valuable comments that improved the discussion. We thank K. Arnaud and G. Madejski, and T. Dotani for help in planning the ASCA observations; A. Vikhlinin, D. Liedahl, G. Mackie, and K. Macleod for useful discussions. This work was supported by NASA grants NAG5–2639 (ASCA), NAG5–1536 and NAG5–2640 (ROSAT), and NASA contract NAS8–39073 (ASC).

REFERENCES

- ABC Guide to ASCA Analysis, http://legacy.gsfc.nasa.gov/docs/asca/abc_contents.html
- COSMOS/UKST Southern Sky Catalogue, <http://wwwpks.atnf.csiro.au/databases/cosmos/COSMOS.html>
- Arnaud K.A., 1996, Astronomical Data Analysis Software and Systems V, eds. Jacoby G. and Barnes J., p17, ASP Conf. Series volume 101.
- Baldwin, J.A., Phillips, M.M., & Terlevich, R. 1981, PASP, 93, 5.
- Boldt E. 1987, Phys Rep, 146, 215
- Boyle, B.J., McMahon, R.G., Wilkes, B.J., & Elvis, M. 1995, MNRAS, 276, 315
- Charles, P.A., & Phillips, M.M. 1982, MNRAS, 200, 263
- Ciliegi P., et al., 1997, MNRAS, 284, 401
- Comastri, A., Setti, G., Zamorani, G., Hasinger, G. 1995, A&A, 296, 1
- Crawford, C.S., & Fabian, A.C. 1994, MNRAS, 267, 779

- David, L.P., Harnden, F.R. Jr., Kearns, K.E., & Zombeck, M.V. 1996 “The ROSAT High Resolution Imager (HRI) Calibration Report,” U.S. ROSAT Science Data Center, Smithsonian Astrophysical Observatory, ftp://legacy.gsfc.nasa.gov/rosat/doc/hri/hri_report/hri.ps.Z.
- Elvis M., Briel U., & Henry J.P., 1983, *ApJ*, 268, 105.
- Elvis, M., Wilkes, B.J., & Lockman, F.J. 1989, *AJ*, 97, 777
- Elvis, M., Fasnacht, C., Wilson, A.S., & Briel, U. 1990, *ApJ*, 361, 459
- Fabian, A.C., Barcons, X., Almaini, O., & Iwasawa, K. 1998, *MNRAS*, in press.
- Fabbiano, G. 1989, *ARA&A*, 27, 87
- Fiore, F. 1997, private communication
- Gendreau, K., Yaqoob, T., Mushotzky, R., & Fabian, A. 1995a, *BAAS*, 187, 5007
- Georgantopoulos, I., Stewart, G.C., Shanks, T., Boyle, B.J., & Griffiths, R.E. 1996, *MNRAS*, 280, 276
- Georgantopoulos I., et al., 1997, *MNRAS*, 291, 203
- Hasinger, G., Burg, R., Giacconi, R., Truemper, J., & Zamorani, G. 1998, *A&A*, 329, 482
- Jones, L.R., McHardy, I.M., Merrifield, M.R., Mason, K.O., Smith, P.J., Abraham, R.G., Branduardi-Raymont, G., Newsam, A.M., Romero-Colmenero, E. 1996 in in *Proc. “Roentgenstrahlung from the Universe,”* eds. Zimmermann, H.U.; Truemper, J.; and Yorke, H.; MPE Report 263, p. 457
- Moorwood, A. F. M. & Oliva, E. 1990, *A&A*, 239, 78
- Morris, S., Ward, M., Whittle, M., Wilson, A.S., Taylor, K. 1985, *MNRAS*, 216, 193
- Morse, J.A. 1994, *PASP*, 106, 67
- Morse, J. A., Wilson, A. S., Elvis, M., & Weaver, K. A. 1995, *ApJ*, 439, 121
- Mushotzky, R.F. 1982, *ApJ*, 256, 92
- Piccinotti, G., Mushotzky, R.F., Boldt, E.A., Holt, S.S., Marshall, F.E., Serlemitsos, P.J., Shafer, R.A. 1982, *ApJ*, 253, 485
- Romero-Colmenero E., Branduardi-Raymont G., Carrera F.J., Jones L.R., Mason K.O., McHardy I.M. & Mittaz J.P.D., 1996, *MNRAS*, 282, 94.
- Schachter, J. F., Remillard, R., Saar, S. H., Favata, F., Sciortino, S., Barbera, M. 1996, *ApJ*, 463, 747
- Schachter, J. F., 1998, in preparation.
- Stocke, J.T. 1995, private communication.
- Stocke, J.T., Morris, S.L., Gioia, I.M., Maccacaro, T., Schild, R., Wolter, A., Fleming, T.A., & Henry, J.P. 1991, *ApJS*, 76, 813
- Turner, T.J. & Pounds, K.A., 1989, *MNRAS*, 240, 833
- de Vaucouleurs, G. H., de Vaucouleurs, A., Corwin, H. G. Jr., Buta, R. J., Paturel, G. & Fouqué, P. 1991, *Third Reference Catalogue of Bright Galaxies* (New York: Springer-Verlag)
- Ward, M.J., Wilson, A.S., Penston, M.V., Elvis, M., Maccacaro, T., & Tritton, K.P. 1978, *ApJ*, 223, 788
- Ward, M., Penston, M.V., Blades, J.C., Turtle, A.J. 1980, *MNRAS*, 193, 563
- Warwick, R.S., Sembay, S., Yaqoob, T., Makishima, K., Ohashi, T., Tashiro, M., & Kohmura, Y. 1993, *MNRAS*, 265, 412

- Weaver, K.A., Yaqoob, T.,
Holt, S.S., Mushotzky, R.F., Matsuoka, M.,
& Yamauchi, M. 1994, ApJ, 436
- Whittle, M. 1992, ApJS, 79, 49
- Wilson, A.S., Elvis, M., Lawrence, A., & Bland-
Hawthorn, J. 1992, ApJ, 391, 75
- Wilson, A. S., Morse, J. A., Elvis, M., Mulchaey,
J. S., Weaver, K. A., Cecil, G. N., & Gal-
limore, J. 1996, in Proc. “Roentgenstrahlung
from the Universe,” eds. Zimmermann, H.U.;
Truemper, J.; and Yorke, H.; MPE Report
263, p. 529
- Wilson, A.S., Nath, B. 1990, ApJS, 74, 731
- Wilson, A.S. 1988, A&A, 206, 41
- Yaqoob T. et al., 1993, MNRAS, 262, 435.

Table 1: Fits to NGC 7582 Combined ASCA SIS0+SIS1 X-ray Spectra^a

Model	χ^2/dof	High State ^b		χ^2/dof	Low State ^b	
		N_H^c	$\Gamma/kT, \Gamma_{unabs}, f_{cov}$		N_H^c	$\Gamma/kT, \Gamma_{unabs}, f_{cov}$
A. Single Power-Law (PL)	233/147	< 0.10	-0.79 ^{+0.09} _{-0.11}	250/146	<0.04	-0.63 ^{+0.10} _{-0.11}
B. PL + Brems. ^d	145/144	4.9 ^{+2.6} _{-1.2}	0.61 ^{+0.50} _{-0.35} /(0.81 ^{+0.90} _{-0.28})	136/143	5.5 $\pm$ 1.3	0.83 ^{+0.43} _{-0.27} /(0.84 ^{+0.46} _{-0.22})
C. Two PLs ^d	146/144	5.3 ^{+2.3} _{-1.2}	0.64 ^{+0.56} _{-0.29} /(2.5 ^{+0.48} _{-0.63})	137/143	6.0 ^{+2.5} _{-1.83}	0.87 ^{+0.59} _{-0.34} /(2.4 ^{+0.34} _{-0.39})
D. Partially Covered PL	158/146	7.9 ^{+1.3} _{-1.2}	1.5 ^{+0.37} _{-0.40} /(0.96 ^{+0.02} _{-0.03})	152/145	9.1 ^{+1.4} _{-1.1}	1.9 ^{+0.32} _{-0.29} /(0.97 $\pm$ 0.01)

Notes: Photon power law indices are used throughout. (a) Ranges of parameters are for 90% confidence level; (b) High and low states are as defined in section 2.1; (c) in units of 10^{22}cm^{-2} ; (d) bremsstrahlung and second power-law absorption fixed to $N_H=N_H(\text{Gal})$.